

PATENT
22851-032

PATENT APPLICATION IN THE U.S. PATENT AND TRADEMARK OFFICE
for
METHOD FOR THE TREATMENT AND PREVENTION OF DENTAL CARIES
by

WENYUAN SHI
MAXWELL H. ANDERSON
SHERIE L. MORRISON
RYAN TRINH
LETITIA WIMS
LI CHEN

Background of the Invention

This application is a continuation-in-part of United States patent application serial number 09/378,577 filed August 20, 1999.

This application relates to an immunologic methodology for the treatment and prevention of dental caries. This invention has special application to patients who are without the ability or motivation to apply established principles of self care, such as very young children, the infirm and poorly educated populations.

Dental caries (tooth decay) and periodontal disease are probably the most common chronic diseases in the world. The occurrence of cavities in teeth is the result of bacterial infection. Hence the occurrence of dental caries is properly viewed as an infectious microbiological disease that results in localized destruction of the calcified tissues of the teeth.

Streptococcus mutans is believed to be the principal cause of tooth decay in man. When *S. mutans* occurs in large numbers in dental plaque, and metabolizes complex sugars, the resulting organic acids cause demineralization of the tooth surface. The result is carious lesions, commonly known as cavities. Other organisms, such as *Lactobaccilli* and *Actinomyces* are also believed to be involved in the progression and formation of carious lesions. Those organisms that cause tooth decay are referred to herein as "cariogenic organisms."

Removal of the damaged portion of a tooth and restoration by filling can, at least temporarily, halt the damage caused by oral infection with cariogenic organisms. However, the "drill and fill" approach does not eliminate the causative bacterial agent. Proper oral hygiene can control the accumulation of dental plaque, where cariogenic organisms grow and attack the tooth surfaces. However, dental self-care has its limits, particularly in populations that are unable to care for themselves, or where there is a lack of knowledge of proper methods of self care. Administration of fluoride ion has been shown to decrease, but not eliminate the incidence of dental caries.

In view of the overwhelming evidence of the involvement of cariogenic organisms in the pathogenesis of dental caries, it is not surprising that there have been a number of different attempts to ameliorate the condition using traditional methods of anti-microbial therapy. The disadvantage of antimicrobial agents is that they are not selective for cariogenic organisms. Administration of non-specific bacteriocidal agents disturbs the balance of organisms that normally inhabit the oral cavity, with consequences that cannot be predicted, but may include creation of an environment that provides opportunities for pathogenic organisms. In addition, long term use of antimicrobial agents is known to select for organisms that are resistant to them. Hence long term and population-wide use of antimicrobial agents to prevent tooth decay is not practical.

Vaccination of humans to elicit an active immune response to *S. mutans*, or other cariogenic organisms, is also not a practical solution at this time. One drawback of this approach is that vaccination elicits production of predominantly IgG and IgM antibodies, but they are not secreted into saliva. The majority of antibodies present in saliva are of the IgA isotype, which can bind to, but cannot activate lymphocytes or complement components to kill bacteria. Accordingly, vaccination is not believed likely to be capable of producing antibodies that can trigger the immune system to kill cariogenic organisms in the mouth. There is no known method for selectively increasing the titer of vaccination induced antibodies of the IgG or IgM isotypes in the oral cavity.

There have been a number of reported attempts to passively immunize patients to *S. mutans* using monoclonal IgA antibodies raised in mice to prevent tooth decay in animals and in man. Because IgA is a multivalent antibody, a single molecule of IgA can bind to several different antigenic sites, resulting in clumping of bacteria. However, binding of IgA to bacterial surface antigens does not kill the bacteria. Rather, clumping is believed to hinder the ability of bacteria to bind to tooth surfaces. Another drawback of this approach is that repeated administration of murine (i.e., heterologous) antibodies to humans has the potential to evoke an immune response to the antibodies.

Unlike IgA antibodies, antibodies of the IgG and IgM classes have bacteriocidal effects. Binding of IgM or IgG antibodies to antigens present on the surface of cariogenic organisms may result in the destruction of the bacterial cells by either of two presently known separate mechanisms: complement mediated cell lysis and antibody-dependent cell-mediated cytotoxicity. In either case, antibodies that selectively bind to certain microbial organisms target just those cells for destruction by the immune system. Both complement mediated cell lysis and antibody-dependent

cell mediated cytotoxicity are part of the humoral immune response that is mediated by antibodies of the IgG and IgM classes.

In order to elicit the desired cytotoxic effect of antibody binding, monoclonal antibodies to cariogenic organisms must be recognized by the human immune system. There are a number of different technologies by which antibodies that will trigger a response from the human immune system can be produced. One example is producing a chimeric antibody using a nucleic acid construct that codes for expression of a human antibody modified to incorporate sequences encoding the variable domain from a different source. Another method utilizes a phage display to determine the actual binding sites of the monoclonal antibody, the complementarity determining regions (CDRs), and then grafting the CDRs onto the framework regions of the variable domains of a human immunoglobulin by site directed mutagenesis. Still other methods, known in the art, which allow the production of antibodies capable of engaging the humoral immune systems include: 1) Immunizing mice which have been genetically altered to produce human antibodies; 2) Immunizing isolated human B cells in vitro and then going through a cell fusion procedure to produce a hybridoma that secretes the antibody; and 3) Isolating B cells from humans with acute infection and producing an antibody generating hybridoma.

Production and administration of such genetically engineered humanized or human monoclonal antibodies to treat dental caries in man poses issues requiring innovative solutions. Prior art methods for production of monoclonal antibodies involve growing hybridomas in culture media, followed by extraction and purification of the desired antibody. These steps are significantly simplified in a preferred embodiment of the invention by expressing the antibodies in edible plants or animals. The antibodies are administered upon oral ingestion of plant or animal products, such as fruits, vegetables or milk wherein the antibodies are not denatured.

This mode of administration has the potential for obviating compliance issues in ameliorating tooth decay.

United States patent application Serial No. 09/378,247 discloses three murine monoclonal antibodies specific to *S. mutans*: SWLA1, SWLA2 and SWLA3. Development of an effective immunological method for the treatment and prevention of dental caries requires preparation of monoclonal antibodies genetically engineered to both express monoclonal antibodies specific to *S. mutans* and engage the effector apparatus of the human immune system.

Summary of the Preferred Embodiments

Dental caries may be prevented or treated by oral ingestion of human or humanized murine monoclonal IgG and IgM antibodies that bind to surface antigens of cariogenic organisms, such as *S. mutans*. The genetically engineered monoclonal antibodies engage the effector apparatus of the human immune system when they bind to cariogenic organisms, resulting in their destruction. In a preferred embodiment, monoclonal antibodies to cariogenic organisms are produced by edible plants, including fruits and vegetables, transformed by DNA sequences that code for expression of the desired antibodies. The genetically engineered monoclonal antibodies are applied by eating the transformed plants.

We have now isolated and sequenced the nucleotide sequences encoding the variable regions of monoclonal antibodies specific to *S. mutans*. When expressed, monoclonal antibodies encoded by these sequences bind specifically to *S. mutans*. Through the use of recombinant techniques, the variable regions of the monoclonal antibodies have been linked to the constant region of human antibodies thereby generating a chimeric monoclonal antibody that specifically binds *S. mutans*. This chimeric monoclonal antibody is directed specifically to surface antigens of

cariogenic organisms which generates an effector response from the immune system upon binding to the target organism.

5

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

10

20

25

Detailed Description of the Preferred Embodiments

1. Preparation of Monoclonal Antibodies

5 The monoclonal antibody technique permits preparation of antibodies with extraordinary specificity. Monoclonal antibodies that bind to specific molecular structures can be produced using what are today considered standard techniques.

10 The monoclonal antibodies that may be used in this invention are those that are directed to surface antigens of cariogenic organisms. Surface antigens are substances that are displayed on the surface of cells. Such antigens are accessible to antibodies present in body fluids. In the context of the present invention, surface antigens of cariogenic organisms are present on the surface of organisms that cause dental caries. While the role of bacterial activity in the genesis of carious lesions is well defined, establishing a cause and effect relationship between a particular
15 organism and the occurrence of dental caries has not been completely successful. To date, only *S. mutans* has been definitively associated with dental caries. However, species of the *Lactobaccili* and *Actinomyces* are also believed to be involved, particularly with the active progression of carious lesions. Any organism that can produce a carious lesion is a potential target for the monoclonal antibodies prepared and used in accordance with this invention.

20 A further requirement of the monoclonal antibodies that may be used in the practice of the present invention is that they are selective for cariogenic organisms. Monoclonal antibodies directed to antigens present on cariogenic as well as non-cariogenic organisms may produce non-specific alterations in the makeup of the flora within the oral cavity. The consequences of such changes are not
25 understood.

Accordingly, the preferred monoclonal antibodies selectively bind to surface antigens of cariogenic organisms. That is to say, the preferred monoclonal antibodies bind specifically to organisms that cause dental caries.

It should be clearly understood that the scope of the present invention is not limited to the prevention of tooth decay in man. Monoclonal antibodies in accordance with the present invention can be genetically engineered to engage the effector response of the immune system of other mammals, such as those that are domesticated as pets.

Monoclonal antibodies can be prepared by immunizing mice or other mammalian hosts with cell wall material isolated from cariogenic organisms. In a preferred embodiment, the cariogenic organisms are type c *S. mutans* (ATCC25175). The immunogenicity of molecules present in cell walls may be enhanced by a variety of techniques known in the art. In a preferred embodiment, immunogenicity of such molecules is enhanced by denaturation of the isolated cell material with formalin. Other techniques for modifying cell wall proteins to enhance immunogenicity are within the scope of this invention. Typically, hosts receive one or more subsequent injections of isolated bacterial cell fragments to increase the titer of antibodies prior to sacrifice and cloning.

Spleen cells from hosts are harvested. The NSI/Ag4.1 mouse myeloma cell line was used as the fusion partner and grown in spinner cultures in 5% CO₂ at 37° C and maintained in log phase of growth prior to fusion. Hybridomas were produced according to the procedure reported by Kohler et al. *Nature*, 256:495-497, (1975). Hybrids were selected in media containing HAT (100 µg Hypoxanthine, 0.4 µM Aminopterin; 16 µM Thymidine). HT (100 µg Hypoxanthine; 16 µM Thymidine) was maintained in the culture medium for 2 weeks after aminopterin was withdrawn. OPI (1 mM oxaloacetate, 0.45 mM pyruvate and 0.2 U/ml bovine insulin) was added as additional growth factors to the tissue culture during cloning of the

hybridomas. The hybridomas were further cloned by limiting dilution using techniques that have become standard since the pioneering work of Kohler and Milstein. In a preferred embodiment, surviving hybridomas were screened for antibody directed to cariogenic organisms by ELISA assay against microtiter plates coated with formalinized bacterial cell material. Positive supernatants were subjected to further screening to identify clones that secrete antibodies with the greatest affinity for the cariogenic organisms. In a preferred embodiment, clones with titers at least three times higher than background are screened again using immunoprecipitation with denatured cell wall material from *S. mutans*. In a preferred embodiment, three clones were identified which bound detectably only to *S. mutans* strains ATCC25175, LM7, OMZ175 and ATCC31377. These clones were deposited with the American Type Culture Collection, receiving Deposit Numbers HB 12599 (SWLA1), HB 12560 (SWLA2), and HB 12558 (SWLA3). United States patent application serial number 09/378,247.

There are various ways to obtain nucleic acid sequences that code for expression of human or humanized monoclonal antibodies specific for the surface antigens of cariogenic organisms: 1) Isolating murine hybridomas which produce monoclonal antibodies against cariogenic organisms and cloning murine genes that code for expression of those antibodies; 2) Using purified cariogenic organisms to screen a phage display random library made from human B lymphocytes to obtain genes that encode antibodies specific for cariogenic organisms; 3) Isolating human hybridomas that produce monoclonal antibodies against cariogenic organisms, using B lymphocytes recovered from heavily infected patients and cloning the human genes encoding these antibodies; or 4) Immunizing human B lymphocytes and spleen cells *in vitro* using purified cariogenic organisms, followed by fusion to form hybridomas to create immortal cell lines. The techniques required are known to those skilled in the art and are not limited to the methods described herein.

2. Preparation of Monoclonal Antibodies Capable Of Eliciting An Effector Response From Human Immune System

Previous efforts to develop an immunological method for the prevention of dental caries employed heterologous antibodies. For example, Lehner, United States patent 5,352,446, refers to use of monoclonal antibodies to *S. mutans* surface antigens raised in mice in inhibiting the proliferation of those bacteria in monkeys. More recently, Ma et al. *Nature Medicine*, 45(5) 601-6 (1998), reported similar results in humans, using a genetically engineered secretory monoclonal murine antibody to *S. mutans* expressed in tobacco plants. Drawbacks to this approach include 1) administration may aggregate the offending organisms, but not kill them because the non-human antibodies do not effectively engage the human immune response; and 2) repeated administration of the antibody may elicit an immune response from the patient to the antibody. A preferable approach is to use recombinant techniques to prepare chimeric antibody molecules directed specifically to surface antigens of cariogenic organisms, that will also elicit an effector response from the immune system of the mammal treated therewith upon binding to the target organism. This can be accomplished by inserting variable regions from murine monoclonal antibodies that are specific to cariogenic organisms into antibodies of the IgG and/or IgM classes from the mammal to be treated. It is also possible to generate antibodies that utilize just the complementarity determining regions (CDRs) of a murine monoclonal antibody specific to cariogenic organisms. Through known recombinant techniques, the CDRs are transferred into the immunoglobulin's variable domain.

Methods are also known for generating the antibody directly, for example: 1) Immunizing mice which have been genetically altered to produce human antibodies; 2) Immunizing isolated human B lymphocytes in vitro and then going through a cell fusion procedure to produce a hybridoma that secretes the antibody; and

3) Isolating B lymphocytes from humans with acute infection and producing an antibody generating hybridoma. The techniques required are known to those skilled in the art. Because each method produces a human antibody, the antibodies are capable of engaging the humoral immune effector systems upon binding to their specific antigens.

5 In the presently preferred embodiment of the invention, chimeric antibodies specific to *S. mutans* were generated. Using PCR or Southern blot techniques, DNA fragments encoding the variable domains of murine hybridomas secreting antibody specific to cell surface antigens of cariogenic organisms were isolated. Using gene cloning techniques, the variable regions were joined to the constant regions of human immunoglobulins. The result of this genetic engineering is a chimeric antibody molecule with variable domains that selectively bind to surface antigens of cariogenic organisms, but which interacts with the human immune effector systems through its constant regions.

10
15 3. Administration of Monoclonal Antibodies

In order to prepare a sufficient quantity of monoclonal antibodies for clinical use, the desired cell line, transfected with sequences encoding the immunoglobulin, must be propagated. Existing technology permits large scale propagation of monoclonal antibodies in tissue culture. The transfected cell lines secrete monoclonal antibodies into the tissue culture medium. The secreted monoclonal antibodies were recovered and purified by gel filtration and related techniques of protein chemistry.

20 In experimental studies, monoclonal antibodies to *S. mutans* have been applied directly to the surface of teeth. Application by ingestion of mouthwash, or by chewing gum has also been proposed. A presently preferred alternative is to express the chimeric monoclonal antibodies of the present invention in edible plants, such as
25 banana or broccoli. Eating plants transformed in accordance with this invention will

result in application of the antibodies to cariogenic organisms present on tooth surfaces, and elsewhere in the mouth. It is also contemplated that other organisms, both plant and animal, may be transformed to express the monoclonal antibodies described herein, so that such antibodies may be ingested, for example, by drinking milk.

5

10
15
20
25

20

25

Brief Description of the Drawings

The present invention is now described, by the way of illustration only, in the following examples which refer to the accompanying FIGS. 1-8, in which:

FIG. 1 shows the DNA sequences (SEQ ID NOS: 1 and 3) encoding the variable regions of the chimeric antibody (TEDW) specific to *S. mutans* derived from SWLA1 cells together with the predicted amino acid sequences (SEQ ID NOS: 2 and 4).

FIG. 2 shows the DNA sequences (SEQ ID NOS: 5 and 7) encoding the variable regions of the chimeric antibody (TEFE) specific to *S. mutans* derived from SWLA2 cells together with the predicted amino acid sequences (SEQ ID NOS: 6 and 8).

FIG. 3 shows the DNA sequences (SEQ ID NOS: 9 and 11) encoding the variable regions of the chimeric antibody (TEFC) specific to *S. mutans* derived from SWLA3 cells together with the predicted amino acid sequences (SEQ ID NOS: 10 and 12).

FIG. 4 shows the DNA sequence (SEQ ID NO: 13) encoding an aberrant light chain variable region derived from SWLA1 cells together with the predicted amino acid sequence (SEQ ID NO: 14).

FIG. 5 shows the DNA sequence (SEQ ID NO: 15) encoding a non-effective heavy chain variable region derived from SWLA1 cells together with the predicted amino acid sequence; (SEQ ID NO: 16).

FIG. 6 shows the DNA sequence (SEQ ID NO: 17) encoding an aberrant heavy chain variable region derived from SWLA1 cells together with the predicted amino acid sequence; (SEQ ID NO: 18).

FIG. 7 shows the DNA sequence (SEQ ID NO: 19) encoding an aberrant heavy chain variable region derived from SWLA2 cells together with the predicted amino acid sequence; (SEQ ID NO: 20).

FIG. 8 shows light and florescent microscope images of chimeric antibody TEDW binding to *S. mutans*.

5

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
220

Examples

1. Producing murine monoclonal antibodies against *S. mutans*

Type c *S. mutans* strain ATCC25175 were grown to log phase in BHI medium and washed twice with phosphate buffered saline, pH 7.2 (PBS), by centrifugation at 3000xg for 5 min. The pellet was resuspended in 1% formalin/0.9% NaCl, mixed at room temperature for 30 min and washed twice with 0.9% NaCl. BALB/c mice (8-10 weeks) were immunized intraperitoneally with 100 μ l of the antigen containing approximately 10^8 whole cells of formalinized intact *S. mutans* bacteria emulsified with Freund's incomplete adjuvant (FIA). After 3-5 weeks, mice received a second dose of antigen (10^8 whole cells of bacteria in FIA). Three days prior to sacrifice, the mice were boosted intravenously with 10^8 whole cells of bacteria in saline.

Spleen cells from hosts were harvested. The tissue culture medium used was RPMI 1640 (Gibco) medium supplemented with 2 mM L-glutamine, 1mM sodium pyruvate, and 10 mM HEPES and containing 100 μ g/ml penicillin and 100 μ g/ml streptomycin with 10% fetal calf serum. The NSI/Ag4.1 mouse myeloma cell line was used as the fusion partner and grown in spinner cultures in 5% CO₂ at 37° C and maintained in log phase of growth prior to fusion. Hybridomas were produced according to the procedure reported by Kohler et al. *Nature*, 256:495-497, (1975). Hybrids were selected in medium containing HAT (100 μ g Hypoxanthine, 0.4 μ M Aminopterin; 16 μ M Thymidine). HT (100 μ g Hypoxanthine; 16 μ M Thymidine) was maintained in the culture medium for 2 weeks after aminopterin was withdrawn. OPI (1 mM oxaloacetate, 0.45 mM pyruvate and 0.2 U/ml bovine insulin) was added as additional growth factors to the tissue culture during cloning of the hybridomas. The hybridomas were further cloned by limiting dilution using techniques that have become standard since the pioneering work of Kohler and Milstein.

The following approach was used for screening for species-specific monoclonal antibodies against *S. mutans*. The initial screening was performed using an ELISA assay, which selects for the culture supernatants containing antibodies that bind to *S. mutans*. Formalinized bacteria were diluted in PBS to $OD_{600} = 0.5$, and added to duplicate wells (100 μ l) in 96 well PVC ELISA plates preincubated for 4 h with 100 μ l of 0.02 mg/ml Poly-L-lysine. These antigen-coated plates were incubated overnight at 4°C in a moist box then washed 3 times with PBS and blocked with 0.5% fetal calf serum in PBS and stored at 4°C. 100 μ l of mature hybridoma supernatants were added to the appropriate wells of the antigen plates, incubated for 1 h at room temperature, washed 3 times with PBS-0.05% Tween 20, and bound antibody was detected by the addition of polyvalent goat-anti-mouse IgG antibody conjugated with alkaline phosphatase diluted 1:1000 with PBS-1% fetal calf serum. After the addition of the substrate, 1 mg/ml p-nitrophenyl phosphate in carbonate buffer (15 mM Na_2CO_3 , 35 mM NaH_2CO_3 , 10 mM $MgCl_2$ pH 9.6), the color development after 15 min was measured in an EIA reader at 405 nm. The positive supernatants (3 fold higher than control) were then subjected to the immunoprecipitation assay (mixing 100 μ l bacteria with 100 μ l supernatant) to screen for those with strong positive reactivity with *S. mutans*. The deposited clones (ATCC HB 12599, HB 12560, and HB 12558) were prepared according to this method.

2. Preparation of Hybridoma lines for cloning of V regions.

A. Isotyping

The hybridoma supernatants were isotyped with a Pharmigen Isotyping Kit (BD Pharmigen, San Deigo, CA). 200 μ l of isotype specific rat anti-mouse antibody was diluted in 800 μ l of coating buffer and 50 μ l of each reagent was added to 10 wells of a 96 well polystyrene ELISA plate. Plates were incubated overnight at 4°C. The plate was washed four times with washing buffer, 0.05% Tween-20 in PBS, and the remaining contents shaken out and the plate blotted dry on a paper towel. 200

5 μ l of blocking solution, 1% BSA in PBS, was added to each well and the plate was incubated at room temperature for 30 minutes. Plates were again washed four times and the contents shaken out. 50 μ l of hybridoma supernatant was added to the appropriate wells. Positive controls from the kit were added to the appropriate wells. The plate was incubated at room temperature for one hour. The plate was washed five times with washing buffer and the plate was blotted dry. One phosphatase substrate tablet was dissolved in 5.0 ml of p-NNP substrate diluent. 50 μ l of substrate solution was added to each well and the plate was incubated for 40 minutes. The plate was read at 405nm on a Dynatech MR 700 microplate reader. SWLA1, SWLA2 and SWLA3 were all determined to be γ 2a, κ (IgG).

B. Biosynthetic labeling

Cells were washed twice in methionine-free, Dulbecco's modified Eagle's medium (DME, Irvine Scientific) supplemented with non-essential amino acids (Grand Island Biological) and glutamine (29.2 μ g/ml). Cells were labeled in 1 ml of DME with 15 μ Ci [35 S]methionine (Amersham, Arlington Heights, IL). All labels were done using 3×10^6 cells.

For labeling of secretions, 35 S-methionine was added to 15 μ C/ml and cells were labeled for 3 hours at 37°C. Cells were harvested on to ice and pelleted by centrifugation. To isolate secreted IgG, the radioactive medium was transferred to a clean tube.

20 For measurement of cytoplasmic IgG, the cell pellet was lysed in 0.5 ml of NDET (1 % NP40, 0.4% deoxycholate, 66mM EDTA and 10mM Tris, pH 7.4), nuclei were pelleted by centrifugation, and the cytoplasmic lysate transferred to a fresh tube. To immunoprecipitate the secreted or cytoplasmic IgG, rat anti-mouse kappa sepharose (prepared in the laboratory) was added. The samples were mixed
25 overnight at 4°C, washed in NDET and then washed with dH₂O. The precipitates were resuspended in sample buffer (25mM Tris, pH 6.7, 2% SDS, 10% glycerol,

0.008% bromophenol blue), and the antibodies were eluted from the sepharose by boiling. The samples were analyzed by SDS-PAGE and autoradiography without reduction on 5% phosphate gels; samples treated with 2-mercaptoethanol were analyzed using 12% tris-glycine gels. Results indicate that all three clones made the same size heavy chain but different size light chains. All three hybridomas were subcloned to ensure homogenous cell populations.

C. Subcloning

The hybridomas were subcloned on soft agar. A 60mm petri dish was coated with 5 ml of growth media plus 10% J774.2 (a murine macrophage cell line) supernatant plus 0.24% agarose (Sigma). The agarose was allowed to harden and a single cell suspension of hybridoma cells mixed with agarose was layered on top. When colonies were about 64 cells in size, they were overlaid with rabbit anti-mouse $\gamma 2a$ specific antiserum mixed with agarose. An immune precipitate forms over and partially obscures those clones secreting $\gamma 2a, \kappa$ antibody. Colonies making the most antibody, were identified and moved up to bulk culture where they were once again biosynthetically labeled.

3. Cloning Variable Regions from SWLA Cells

The basic protocol for cloning the variable regions from SWLA cells is outlined below followed by its application to specific SWLA hybridomas.

- (i) Murine mRNA is made from about 5×10^6 of both the original and subcloned cells using the Microfast Track Kit from Invitrogen.
- (ii) First strand cDNA is made using oligonucleotides that prime the 5' of the light or heavy chain constant region or that prime to the polyA tail of mRNA.

Basic Protocol:

- 5
- (a) Half of the cDNA is resuspended in 20 μ l RNase free dH₂O
 - (b) 2 μ l of 0.5 mg/ml primer is added; the mixture is incubated @ 60°C for 10 minutes
 - (c) The sample is cooled on ice; 8 μ l of 5X first strand cDNA buffer, 2 μ l of RNasin (Promega), 4 μ l of 5mM dNTP, and 0.5 μ l of AMV Reverse Transcriptase are then added
 - (d) The sample is incubated @ 42°C for 1 hour
- 10
- (iii) PCR amplification is done with a number of different light or heavy chain signal peptide primers and primers that hybridize 5' of the light or heavy chain constant region.

PCR Conditions:

- 15
- (a) Denature @ 94°C for 40 sec.
 - (b) Anneal @ 60°C for 40 sec.
 - (c) Extend @ 72°C for 40 sec.
 - (d) Amplify for 30 cycles
 - (e) Final Extension at 72°C for 10 min.
- 20
- (iv) The resulting PCR products are cloned into Invitrogen's PCR2.1 vector via the TOPO Cloning Kit.
 - (v) Individual clones were sent out for sequencing. The results were analyzed for an open reading frame (ORF) and compared with the known database to ensure that the sequence cloned is a variable region.
- 25
- (vi) To check for PCR induced nucleotide alterations in the sequence, steps III to V were repeated so that the sequence of different clones from independent PCR reactions can be compared to ensure the accuracy of the sequence. The

sequence data are also used to determine the sequence of the J region primer that needs to be used.

- (vii) The variable region was cloned into the proper light (human kappa) or heavy chain (human IgG1) expression vector.

Typical Variable Region Ligation Protocol:

- (a) 1 μ g each of vector and insert was cut with either NheI/EcoRV or SalI/EcoRV
- (b) The relevant fragments were then isolated using Qiagen's Gel Extraction Kit
- (c) The ligation reaction used 4 μ l out of 30 μ l of vector sample and 6 μ l out of 30 μ l of insert sample
- (d) 4 to 5 units of T4 DNA ligase was then added for a 20 μ l final reaction volume

Typical Transformation Protocol:

- (a) An aliquot of HB101 chemically competent *E. coli* cells were thawed on ice
- (b) The entire ligation reaction was then added to the cells and incubated on ice for 15 minutes
- (c) The cells were then heat shocked @ 42°C for 1 minute
- (d) 1 ml of LB was added to the cells and the tube was shaken @ 37°C for 1 hour
- (e) The tube was spun down @ 8000 RPM for 2 minutes
- (f) All but 100 μ l of supernatant was removed
- (g) The cells were resuspended, plated onto LB+AMP plates, and incubated @ 37°C overnight

B. Cloning the Variable Regions from SWLA1 cells

In attempting to clone the light chain variable region (VL), PCR product was found using signal peptide primer 442 with constant region primer 450 as shown below. Previous studies have determined that 442 also primes to an endogenous aberrant or non-productive VL, SWLA1 Aberrant VL (SEQ ID NO: 13). Knowing this, attempts were made to enrich for non-aberrant transcripts by restriction digesting the PCR product with PflMI, which recognizes a specific sequence in the aberrant VL. Eventually, one variable region sequence was found to have an ORF. The final PCR product SWLA1 VL (SEQ ID NO: 1) was generated with primer 442 and J region primer 453 as shown below and inserted into the appropriate expression vector. The resulting human kappa expression vector carrying the VL from SWLA1 is named 5936 pAG.

See FIG. 1 Panel A which shows the sequence coding the VL domain and the predicted amino acid sequence (SEQ ID NOS: 1 and 2) and FIG. 4 which shows the sequence coding the aberrant VL and the predicted amino acid sequence (SEQ ID NOS: 13 and 14)

442 (SEQ ID NO: 21) 5' GGG GAT ATC CAC ATG GAG ACA GAC ACA
CTC CTG CTA T 3'

450 (SEQ ID NO: 22) 5' GCG TCT AGA ACT GGA TGG TGG GAA GAT
GG 3'

453 (SEQ ID NO: 23) 5' AGC GTC GAC TTA CGT TTK ATT TCC ARC
TTK GTC CC 3'

The cloning of the heavy chain variable region (VH) resulted in finding two unique VHs both with ORFs. One VH uses signal peptide primer 440 and the other uses signal peptide primer 441 as shown below. In both reactions, the heavy chain constant region primer 451 was used. Two final PCRs were done. The first

used J region primer 452 with primer 440 which generated SWLA1 VH (SEQ ID NO: 3) and the second used the same J region primer with primer 441 which produced SWLA1 2nd VH (SEQ ID NO: 15) and an aberrant non-productive VH, SWLA1 Aberrant VH (SEQ. ID NO: 17). The resulting human IgG1 expression vectors carrying the two different VHs generated are named 5937 pAH (SWLA1 VH) and 5943 pAH (SWLA1 2nd VH). Only vector 5937 pAH however was found to express an effective full length VH.

The DNA coding the VH domain and the predicted amino acid sequence are shown in FIG. 1 Panel B as SEQ ID NOS: 3 and 4. See FIG. 5 for the non-effective 2nd VH DNA and amino acid sequence (SEQ ID NOS: 15 and 16) and FIG. 6 for the DNA and amino acid sequence for the aberrant VH (SEQ ID NOS: 17 and 18).

440 (SEQ ID NO: 24) 5' GGG GAT ATC CAC ATG RAC TTC GGG YTG
AGC TKG GTT TT 3'

441 (SEQ ID NO: 25) 5' GGG GAT ATC CAC ATG GCT GTC TTG GGG
CTG CTC TTC T 3'

451 (SEQ ID NO: 26) 5' AGG TCT AGA AYC TCC ACA CAC AGG RRC
CAG TGG ATA GAC 3'

452 (SEQ ID NO: 27) 5' TGG GTC GAC WGA TGG GGS TGT TGT GCT
AGC TGA GGA GAC 3'

C. Cloning the Variable Regions from SWLA2 cells

Two PCR products were found in cloning the VL. One product came from primers 442 and 450. The other came from primer 443 and primer 450. A unique VL with an ORF was cloned from the 443 and 450 reaction. The final PCR, which generated the SWLA2 VL (SEQ ID NO: 5), used J region primer 453 with

primer 443. The resulting human kappa expression vector carrying the VL from SWLA2 is named 5938 pAG.

See FIG. 2 Panel A which shows the sequence coding the VL domain and the predicted amino acid sequence (SEQ ID NOS: 5 and 6).

443 (SEQ ID NO: 28) 5' GGG GAT ATC CAC ATG GAT TTT CAA GTG
CAG ATT TTC AG 3'

Two PCR products were also found in cloning the VH. One product came from primers 439 and 451. The other product came from primers 440 and 451. The transcript from the former reaction turned out to be aberrant, SWLA2 Aberrant VH (SEQ ID NO: 19). The transcript from the latter reaction was missing part of its 5' sequence. After aligning this sequence to several similar known VHs, a new leader signal peptide primer 843 was designed as shown below. The final PCR product SWLA2 VH (SEQ ID NO: 7) was generated with primer 843 with J region primer 452. The resulting human IgG1 expression vector carrying the VH from SWLA2 is named 5939 pAH.

The DNA coding the VH domain and the predicted amino acid sequence are shown in FIG. 2 Panel B as SEQ ID NOS: 7 and 8. See FIG. 7 for the DNA and amino acid sequence for the aberrant VH (SEQ ID NOS: 19 and 20).

439 (SEQ ID NO: 29) 5' GGG GAT ATC CAC ATG GRA TGS AGC TGK
GTM ATS CTC TT 3'

843 (SEQ ID NO: 30) 5' GGG ATA TCC ACC ATG GRC AGR CTT ACW
TYY TCA TTC CTG 3'

D. Cloning the Variable Regions from SWLA3 cells

The only VL PCR product came from primer combination 442 and 450. Once again the PCR product was digested with PflMI to enrich for non-aberrant transcripts. This procedure didn't help. Another enzyme Eco0109I was used similarly and one transcript was found with the 5' end missing. The sequence was compared to the known database and a new signal peptide primer 826 was designed as shown below. This primer 826 was then used with J region primer 835 shown below to yield the final PCR product SWLA3 VL (SEQ ID NO: 9). It was cloned into a human kappa expression vector and named 5940 pAG.

See FIG. 3 Panel A which shows the sequence coding the VL domain and the predicted amino acid sequence (SEQ ID NOS: 9 and 10).

826 (SEQ ID NO: 31) 5' GGG GAT ATC CAC ATG ATG AGT CCT GCC
CAG TTC C 3'

835 (SEQ ID NO: 32) 5' GGT CGA CTT AGC TTT CAG CTC CAG CTT
GGT 3'

The only VH PCR product was obtained from primer combination 440 and 451. The final PCR reaction used primer 440 and J region primer 452 to generate SWLA3 VH (SEQ ID NO: 11). The VH was cloned into a human IgG1 expression vector and named 5941 pAH.

The DNA coding the VH domain and the predicted amino acid sequence are shown in FIG. 3 Panel B as SEQ ID NOS: 11 and 12.

4. Generating murine/human chimeric genes which encode humanized monoclonal antibodies against *S. mutans*.

- (i) DNA was prepared from the expression vectors and from the plasmid containing the correct V regions. See *Current Protocols in Immunology*, Section 2.12.1 (1994) for detailed information about the vectors that express the light and heavy chain constant regions.

- (ii) The expression vector was digested with the appropriate restriction enzyme. The digests were then electrophoresed on an agarose gel to isolate the appropriate sized fragment.
- (iii) The plasmid containing the cloned V region was also digested and the appropriate DNA fragment containing the V region was isolated from an agarose gel.
- (iv) The V region and expression vector were then mixed together, T4 DNA ligase was added and the reaction mixture was incubated at 16°C over night.
- (v) Competent cells were transfected with the ligation mixture and the clones expressing the correct ligation sequence were selected. Restriction mapping was used to confirm the correct structure.

5. Transfecting eukaryotic cells

10 micrograms of DNA from each expression vector was linearized by BSPC 1 (Stratagene, PvuI isoschizomer) digestion and 1×10^7 myeloma cells (Sp2/0 or NSO/1) were cotransfected by electroporation. Prior to transfection the cells were washed with cold PBS, then resuspended in 0.9 ml of the same cold buffer and placed in a 0.4 cm electrode gap electroporation cuvette. 960 microF and 200V were used for electroporation. The shocked cells were then incubated on ice in IMDM medium (Gibco, NY) with 10% calf serum.

The transfected cells were plated into 96 well plates at a concentration of 10^4 cells/well. Selective medium including selective drugs such as histidinol or mycophenolic acid were used to select the cells which contain expression vectors. After 12 days, the supernatants from growing clones were tested for antibody production.

6. Analyses of recombinant antibodies

ELISA assay was used to identify transfectomas that secrete human IgG antibodies. 100 µl of 5 µg/ml goat anti-human IgG was added to each well of a

96-well ELISA plate and incubated overnight. The plate was washed several times with PBS and blocked with 3% BSA. Supernatants from above growing clones were added to the plate for 2 hours at room temperature. Plates were then washed and anti-human kappa antibody labeled with alkaline phosphatase diluted 1:10,00 in 1% BSA was added for 1 hour at 37° C. Plates were washed with PBS and p-NPP in diethanolamine buffer (9.6% diethanolamine, 0.24 mM MgCl₂, pH 9.8) was added. Color development at OD₄₀₅ was indicative of cells producing H₂L₂.

For the supernatants that produce humanized IgG constant regions, their reactivity with *S. mutans* was tested as described in Shi et al., *Hybridoma* 17:365-371 (1998). Briefly, bacteria strains listed in Table 1 were grown in various media suggested by the American Type Culture Collection. The anaerobic bacteria were grown in an atmosphere of 80% N₂, 10% CO₂, and 10% H₂ at 37° C. The specificity of antibodies to various oral bacteria was assayed with ELISA assays. Bacteria were diluted in PBS to OD₆₀₀=0.5, and added to duplicate wells (100 µl) in 96 well PVC ELISA plates preincubated for 4 h with 100 µl of 0.02 mg/ml Poly-L-lysine. These antigen-coated plates were incubated overnight at 4° C in a moist box then washed 3 times with PBS and blocked with 0.5% fetal calf serum in PBS and stored at 4° C. 100 µl of chimeric antibodies at 50 µg/ml were added to the appropriate wells of the antigen plates, incubated for 1 h at RT, washed 3 times with PBS-0.05% Tween 20, and bound antibody detected by the addition of polyvalent goat-anti-human IgG antibody conjugated with alkaline phosphatase diluted 1:1000 with PBS-1% fetal calf serum. After the addition of the substrate, 1 mg/ml p-nitrophenyl phosphate in carbonate buffer (15 mM Na₂CO₃, 35 mM NaH₂CO₃, 10 mM MgCl₂ pH 9.6), the color development after 15 min was measured in a EIA reader at 405 nm. "+" means OD₄₀₅>1.0; "-" means OD₄₀₅<0.05. The negative control is <0.05. Chimeric antibodies used are TEDW (derived from SWLA1), TEFE

(derived from SWLA2) and TEFC (derived from SWLA3). The results are given in Table 1.

TABLE 1

Reactivity of Chimeric Antibodies to Various Oral Bacterial Strains		
Oral Bacteria	Strains	Chimeric antibodies
<i>S. mutans</i>	AATCC25175	+
	LM7	+
	OMZ175	+
	ATCC49456	-
	ATCC19645	-
	ATCC49295	-
	ATCC6715-B	-
	ATCC33478	-
	ATCC4356	-
	ATCC11578	-
	ATCC14917	-
	ATCC11742	-
	ATCC33384	-
	ATCC12104	-
	ATCC19246	-
	ATCC25586	-
	ATCC33277	-

FIG. 8 shows fluorescent microscopy images generated using the chimeric TEDW antibody derived from SWLA1. *S. mutans* ATCC25175 was grown in Brain-Heart Infusion medium in an atmosphere of 80% N₂, 10% CO₂, and 10% H₂ at 37°C. Bacteria were then washed and resuspended in PBS buffer, mixed with various antibodies and examined with light microscopy or fluorescent microscopy. Referring to FIG. 8: Left, chimeric antibodies bind and agglutinate *S. mutans* cells; middle, chimeric antibodies interact with goat, FITC conjugated anti-human IgG (Fc specific) antibody (Sigma F9512) to give fluorescent image of *S. mutans*; right, chimeric antibodies do not react with goat, FITC conjugated anti-mouse IgG (Fc

specific) antibody (Sigma F5387) and give no fluorescent image of *S. mutans*. Chimeric antibodies TEFE and TEFC were also used and produced results consistent with the TEDW chimeric antibody.

Results from both the flow cytometry and florescent microscopy experiments indicate that each chimeric antibody (TEDW, TEFE, and TEFC) contained both a human IgG constant region and a variable region capable of specifically recognizing *S. mutans*.

7. Expressing Monoclonal Antibodies to *S. mutans* In Transformed Organisms

A. Producing human or humanized monoclonal antibodies in animal cells

The heavy and light chain of a human IgG gene are separately introduced or cotransfected into an animal cell line (such as Sp2/0) using electroporation. The transfected cells are plated onto a microtiter plate and incubated at 37° C in a 5% CO₂ atmosphere in medium containing 10% fetal bovine serum. After a 48 h incubation, the cells are grown in selection medium containing histidinol or mycophenolic acid. The supernatants of drug-resistant cells are collected and screened for immuno-reactivity against *S. mutans* using the ELISA or precipitation assays mentioned above.

B. Producing human or humanized monoclonal antibodies in edible plants

Transgenic plants have been recognized as very useful systems to produce large quantities of foreign proteins at very low cost. Expressing human or humanized monoclonal antibodies against *S. mutans* in edible plants (vegetables or fruits) allows direct application of plant or plant extracts to the mouth to treat existing dental caries and to prevent future bacterial infection. The choice of transgenic, edible plants includes, but is not limited to, potato, tomato, broccoli, corn, and banana.

Presented here are the procedures to produce transgenic *Arabidopsis*, an edible plant closely related to *Brassica* species including common vegetables such as cabbage, cauliflower and broccoli. It is chosen because many genetic and biochemical tools have been well developed for this plant. There are several strategies to express IgG in this plant. One strategy is to first introduce the human IgG genes encoding the heavy chain and light chain to two separate transgenic lines. The two genes are brought together by genetic crossing and selection. Other methods involve sequential transformation, in which transgenic lines transformed with one IgG gene are re-transformed with the second gene. Alternatively, genes encoding the heavy chain and light chain are cloned into two different cloning sites in the same T-DNA transformation vector under the control of two promoters, and the expression of both genes can be achieved by the transformation of a single construct to plant. Technically, the separate transformation method is the simplest one and it usually results in higher antibody yield. Therefore, we present this strategy here. It is possible to transform other plants using similar techniques.

The DNA fragments encoding the heavy and light chains of a human IgG gene are separately cloned into a Ti plasmid of *Agrobacterium tumefaciens*. The plasmid contains a promoter to express human heavy and light chains of IgG in *Arabidopsis thaliana*, an antibiotic marker for selection in *Agrobacterium tumefaciens* and an herbicide resistance gene for transformation selection in *Arabidopsis*. An *Agrobacterium tumefaciens* strain is transformed with these plasmids, grown to late log phase under antibiotic selection, and resuspended in infiltration medium described by Bethtold et al. (C.R. Acad. Sci. Paris Life Sci. 316:1194-1199, 1993).

Transformation of *Arabidopsis* by Ti-plasmid containing *Agrobacterium tumefaciens* is performed through vacuum infiltration. Entire plants of *Arabidopsis* are dipped into the bacterial suspension. The procedure is performed in a vacuum chamber. Four cycles of 5 min vacuum (about 40 cm mercury) are

applied. After each application, the vacuum is released and reapplied immediately. After infiltration, plants are kept horizontally for 24 h in a growth chamber. Thereafter, the plants are grown to maturity and their seeds are harvested. The harvested seeds are germinated under unselective growth condition until the first pair of true leaves emerged. At this stage, plants are sprayed with the herbicide Basta at concentration of 150 mg/l in water. The *aribidopsis* plants containing transformed Ti plasmids are resistant to the herbicide while the untransformed plants are bleached and killed. Such a selection continues to the second generation of the plants. For the resistant plants, total genomic DNA is isolated and probed with the DNA fragments encoding heavy and light chains of the IgG gene. The plant extracts from the positive transformants are prepared and screened for the expression of human IgG protein with Western blot using antibodies against heavy and light chains of constant regions of human IgG.

The plants expressing human IgG heavy chain are sexually crossed with plants expressing human IgG light chain to produce progeny expressing both chains. Western blotting is used to screen the both heavy and light chains. Extracts from positive transformants are collected and screened for immuno-reactivity against *S. mutans* using the ELISA or precipitation assays mentioned above.

8. Using human or humanized monoclonal antibodies against *S. mutans* to treat or prevent human dental caries

With the successful completion of the above studies, humanized monoclonal antibodies against *S. mutans* are obtained. The plant tissue is tested for efficacy.

Plant tissue extracts containing monoclonal antibodies to *S. mutans* are mixed with various concentrations of *S. mutans* in the presence and absence of purified human complement components or purified human polymorphonuclear

neutrophilic leukocytes. After a two hour incubation, the mixtures are plated onto BHI plates to examine the bactericidal activity.

Using the artificial plaque formation system developed by Wolinsky et al., *J. Dent. Res.* 75:816-822 (1996), plant tissue extracts containing monoclonal antibodies are used to examine the ability of the expressed monoclonal antibodies to kill *S. mutans* in saliva or in existing dental plaques on artificial dental enamel. Analogous techniques are used to examine the ability to prevent the formation of dental plaques.

Human clinical trials are performed using these monoclonal antibodies produced through animal cells or plants. Human volunteers are treated with or without these human monoclonal antibodies against *S. mutans*. Then the level of *S. mutans* in saliva and in dental plaques is examined. The correlation between present and future dental caries in relationship with treatment of monoclonal antibodies is also examined.

It should be understood that the foregoing examples are for illustrative purposes only, and are not intended to limit the scope of applicants' invention which is set forth in the claims appearing below.